

# COMMANDER SF

## 8-32-5

### GUARANTEED ANALYSIS

Total Nitrogen (N).....	8.0%
8.0% Ammoniacal Nitrogen	
Available Phosphate (P2O5).....	32.0%
Soluble Potash(K2O).....	5.0%
Zinc(Zn).....	0.05%

**DERIVED FROM:** Ammonium Poly Phosphate, Potassium Tri-Poly Phosphate, Phosphoric Acid, and Zinc Sulfate.

**ALSO CONTAINS NON-PLANT FOOD INGREDIENT:** 0.2% Humic Acid (Derived from Leonardite)

Follow manufacturer's recommended directions for use.

## SHAKE WELL

**KEEP OUT OF REACH OF CHILDREN**

**CAUTION:** Harmful if swallowed. Avoid breathing spray mist. May cause irritation of nose, throat or skin. Avoid contact with eyes, skin or clothing. In case of contact with eyes, flush with copious amounts of water. Consult physician if irritation persists.

**Wt per gal:** 11.81 lb @ 68°F    1.42 Kg/Ltr @ 20°C

TRIANGLE  
**TRICARD**  
CARDINAL

MANUFACTURED FOR:  
Triangle Chemical  
P.O. Box 4528  
Macon, GA  
31208 F1486

## APPLICATION DIRECTIONS

### DIRECTIONS FOR USE

**TURF GRASSES:** Apply at 2-8 oz/1000 sq. ft. in adequate water for sufficient coverage.

### FIELD CROPS:

**Alfalfa (hay):** Apply 2 - 4 quarts per acre 5 days after each cutting.

**Alfalfa (seed):** Apply 3 - 4 quarts at pre-bloom and several more times at 3-week intervals.

**Barley, Wheat, Small Grains:** Apply 2 - 3 quarts per acre when plant is 10 inches, and again prior to heading out.

**Cotton:** Apply 2 quarts per acre at first true leaf. Repeat at first square and again at 2-week intervals.

**Peanuts:** Apply 3-4 quarts per acre in soil as a starter. Apply 1-3 quarts 1-2 weeks prior to bloom. Repeat again 2 weeks after bloom.

**Rice:** Apply 2-3 quarts per acre at boot stage and 1-2 weeks prior to bloom. Repeat again 2 weeks after bloom.

**Tobacco:** Apply 1-2 quarts per acre in transplant water. Apply 1-3 quarts after transplant. Repeat at 2-3 week intervals.

### VEGETABLE CROPS:

**Artichokes, Beans, Peas, Melons, Squash, Cucumbers, Tomatoes, Peppers:** Apply 2 - 4 quarts per acre at bud stage, prior to bloom and then an additional application every 2 weeks.

### IN FURROW APPLICATIONS:

**Corn, Soybeans, Peanuts:** Apply 2 - 4 quarts per acre. Sweet Potatoes: apply 2-4 quarts per acre in Transplant water

**Cole Crops (Transplant and Direct Seed):** Apply 1 - 2 quarts at 4-6 true leaf stage, 2 - 3 quarts at bud stage, and 2 - 3 quarts at two week intervals.

**Corn:** Apply 2 - 3 quarts when plant is 21 inches and again prior to tasseling. **Lettuce, Spinach, Celery:** Apply 2 - 3 quarts after thinning spinach and lettuce. Apply 2 weeks after celery transplant. Apply again every 2 weeks thereafter. **Potatoes, Yams:** Apply 2 - 3 quarts when plants are 6 inches high. Apply two more times at three week intervals.

**Soybeans:** Apply 2 - 3 quarts at bud stage and repeat again at pod-fill.

**Other Vegetable Crops and Root Crops:** Apply 2 to 3 quarts per acre every 2 weeks after plants are established.

### TREE, VINE, BERRY CROPS:

**Apples and Pears:** Use 2 to 4 quarts per acre at pre-bloom, petal fall, and once again 3-4 weeks later. Do not apply to Melrose Apples.

**Apricots, Cherries, Almonds, Plums, Peaches, Prunes, Nectarines:** Apply 2-4 quarts pre-bloom. Repeat rate after petal fall. A third application may be required.

**Citrus:** Use 4 to 6 quarts per acre at early bud stage, prior to bloom, and again at early fruit set and at fruit size. Do not combine with oils.

**Boysenberries, Blackberries, Cranberries Blueberries, Raspberries:** Apply 2 - 3 quarts / acre at pre-bloom and at 2 week intervals thereafter.

**Grapes:** Apply 1-2 quarts per acre when canes reach 15 - 20 inches. Apply another 2 - 3 quarts at pre-bloom buckshot stage and at 4-6 weeks prior to harvest.

**Pecans, Pistachios, Walnuts:** Apply 3-4 quarts per acre at pre-bloom, and again every 4 - 6 weeks until 30 days prior to harvest.

**Strawberries:** Apply 2 - 3 quarts per acre at 2-week intervals beginning at pre-bloom and continuing throughout the season.